

Product datasheet for AP52934PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

NOR1 (NR4A3) (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1/1000.

Western Blot: 1/100 - 1/500.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide between 552-582 amino acids from the C-terminal region

of human NR4A3

Specificity: This antibody reacts to NR4A3.

Formulation: PBS

State: Aff - Purified

State: Liquid purified Ig fraction

Preservative: 0.09% (W/V) sodium azide

Concentration: lot specific

Purification: Affinity chromatography on Protein A

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 68230 Da

Gene Name: nuclear receptor subfamily 4 group A member 3

Database Link: Entrez Gene 8013 Human

Q92570






Background:

This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. The encoded protein may act as a transcriptional activator. The protein can efficiently bind the NGFI-B Response Element (NBRE). Three different versions of extraskeletal myxoid chondrosarcomas (EMCs) are the result of reciprocal translocations between this gene and other genes. The translocation breakpoints are associated with Nuclear Receptor Subfamily 4, Group A, Member 3 (on chromosome 9) and either Ewing Sarcome Breakpoint Region 1 (on chromosome 22), RNA Polymerase II, TATA Box-Binding Protein-Associated Factor, 68-KD (on chromosome 17), or Transcription factor 12 (on chromosome 15). Multiple transcript variants encoding different isoforms have been found for this gene.

Synonyms: Cl

CHN, CSMF, MINOR, NOR1, TEC

Product images:



NR4A3 Antibody (C-term) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the NR4A3 antibody detected the NR4A3 protein (arrow).