

Product datasheet for **AP53022PU-N**

OR1D5 (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1/1000. Western Blot: 1/1000.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	KLH conjugated synthetic peptide between 244-271 amino acids from the C-terminal region of Human OR1D5.
Specificity:	This antibody recognizes Human Olfactory receptor 1D5 (C-term). Other species not tested.
Formulation:	PBS containing 0.09% (W/V) Sodium Azide as preservative State: Aff - Purified State: Liquid purified Ig fraction
Concentration:	lot specific
Purification:	Protein A column, followed by peptide affinity purification
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	olfactory receptor family 1 subfamily D member 5
Database Link:	Entrez Gene 8386 Human P58170



[View online »](#)

Background:

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

Synonyms:


OR1D5

Note:**Molecular Weight:** 35424 Da**Protein Families:**

Druggable Genome, Transmembrane

Protein Pathways:

Olfactory transduction

Product images:

Western blot analysis of OR1D5 Antibody (C-term) pre-incubated without (Lane 1) and with (Lane 2) blocking peptide in human placenta tissue lysate. OR1D5 Antibody (C-term) (arrow) was detected using the purified Pab (1:60/250 dilution).