

## **Product datasheet for AP53198PU-N**

## PCDHA4 (C-term) Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1/1000.

Western blot: 1/100 - 1/500.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

**Immunogen:** KLH conjugated synthetic peptide between 760-790 amino acids from the C-terminal region

of human PCDHA4

**Specificity:** This antibody reacts to PCDHA4.

Formulation: PBS

State: Aff - Purified

State: Liquid purified Ig fraction

Preservative: 0.09% (W/V) sodium azide

**Concentration:** lot specific

**Purification:** Affinity chromatography on Protein A

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 102293 Da

**Gene Name:** protocadherin alpha 4

**Database Link:** Entrez Gene 56144 Human

Q9UN74



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene is a member of the protocadherin alpha gene cluster, one of three related gene clusters tandemly linked on chromosome five that demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The alpha gene cluster is composed of 15 cadherin superfamily genes related to the mouse CNR genes and consists of 13 highly similar and 2 more distantly related coding sequences. The tandem array of 15 N-terminal exons, or variable exons, are followed by downstream C-terminal exons, or constant exons, which are shared by all genes in the cluster. The large, uninterrupted N-terminal exons each encode six cadherin ectodomains while the C-terminal exons encode the cytoplasmic domain. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins that most likely play a critical role in the establishment and function of specific cell-cell connections in the brain. Alternative splicing has been observed and additional variants have been suggested but their full-length nature has yet to be determined.

**Synonyms:** Protocadherin alpha-4, PCDH-alpha-4

**Protein Families:** Transmembrane

## **Product images:**



PCDHA4 Antibody (C-term) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the PCDHA4 antibody detected the PCDHA4 protein (arrow).