

Product datasheet for AP53512PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD45 (PTPRC) (N-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1/1000.

Western Blot: 1/100-1/500.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide between 343-373 amino acids from the N-terminal region

of Human CD45 / LCA.

Specificity: This antibody recognizes Human CD45 / LCA (N-term).

Formulation: PBS

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.09% Sodium Azide

Concentration: lot specific

Purification: Affinity Chromatography on Protein A

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: protein tyrosine phosphatase, receptor type C

Database Link: Entrez Gene 5788 Human

P08575





Background: The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP)

family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus belongs to receptor type PTP. This gene is specifically expressed in hematopoietic cells. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Four alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq].

Synonyms: PTPRC, Leukocyte common antigen, L-CA, T200


Note: Molecular Weight: 147254 Da

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Phosphatase, Transmembrane **Protein Pathways:**

Cell adhesion molecules (CAMs), Fc gamma R-mediated phagocytosis, Primary

immunodeficiency, T cell receptor signaling pathway

Product images:



Western blot analysis of CD45 / LCA Antibody (Nterm) in Jurkat cell line lysates (35ug/lane).