

Product datasheet for **AP54697PU-N**

ZNF384 (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	FC, WB
Recommended Dilution:	ELISA: 1:1;000. Western blot: 1:100~500. Flow cytometry: 1:10~50.
Reactivity:	Human
Host:	Rabbit
Isotype:	Ig
Clonality:	Polyclonal
Immunogen:	KLH conjugated synthetic peptide between 538-567 amino acids from the C-terminal region of human ZNF384
Specificity:	This antibody detects ZNF384 (C-term).
Formulation:	PBS with 0.09% (W/V) sodium azide State: Aff - Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Protein A column followed by peptide affinity purification
Conjugation:	Unconjugated
Storage:	Store at 2 - 8 °C for up to six months or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	zinc finger protein 384
Database Link:	Entrez Gene 171017 Human Q8TF68



[View online »](#)

Background:

This gene contains long CAG trinucleotide repeats coding consecutive glutamine residues. The gene product may function as a transcription factor, with a potential role in the regulation of neurodevelopment or neuroplasticity. The protein appears to bind and regulate the promoters of MMP1, MMP3, MMP7 and COL1A1. Studies in mouse suggest that nuclear matrix transcription factors (NP/NMP4) may be part of a general mechanical pathway that couples cell construction and function during extracellular matrix remodeling. Multiple transcript variants encoding several isoforms have been found for this gene.

Synonyms:


Zinc finger protein 384, CAGH1, CIZ, NMP4, TNRC1

Note:


Molecular Weight: 63219 Da

Protein Families:

Transcription Factors

Product images:


ZNF384 Antibody (C-term) western blot analysis in MCF-7, HeLa cell line lysates (35 ug/lane). This demonstrates the ZNF384 antibody detected the ZNF384 protein (arrow).



ZNF384 Antibody (C-term) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.