

Product datasheet for AP54734PU-N

ZNF98 (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: FC, IHC, WB

Recommended Dilution: ELISA: 1:1;000.

Western blot: 1:100~500.

Immunohistochemistry: 1:50~100.

Flow cytometry: 1:10~50.

Reactivity: Human Host: Rabbit

Isotype: lg

Clonality: Polyclonal

Immunogen: KLH conjugated synthetic peptide between 486-516 amino acids from the C-terminal region

of human ZNF98

Specificity: This antibody detects ZNF98 (C-term).

Formulation: PBS with 0.09% (W/V) sodium azide

State: Aff - Purified State: Liquid Ig fraction

Concentration: lot specific

Purification: Protein A column followed by peptide affinity purification

Conjugation: Unconjugated

Storage: Store at 2 - 8 °C for up to six months or (in aliquots) at -20 °C for longer. Avoid repeated

freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: zinc finger protein 98

Database Link: Entrez Gene 148198 Human

A6NK75

Background: May be involved in transcriptional regulation.

Synonyms: Zinc finger protein 98,ZNF739



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Note: Molecular Weight: 61144 Da

Product images:




ZNF98 Antibody (C-term) western blot analysis in MDA-MB435 cell line lysates (35 ug/lane). This demonstrates the ZNF98 antibody detected the ZNF98 protein (arrow).



ZNF98 antibody (C-term) immunohistochemistry analysis in formalin fixed and paraffin embedded human lymphnode followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZNF98 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



MDA-MB435



ZNF98 Antibody (C-term) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITCconjugated goat-anti-rabbit secondary antibodies were used for the analysis.