

## Product datasheet for **AP55068SU-N**

### c Maf (MAF) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, EMSA, IHC, WB
Recommended Dilution:	<b>ELISA.</b> <b>Western Blot:</b> 1/200-1/5000. <b>Immunohistochemistry:</b> 1/50-1/500. Useful for <b>Electrophoretic Mobility Shift Assay (EMSA).</b>
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Maf-GST fusion protein
Specificity:	This antibody reacts with 47 kDa Maf protein derived from a transfected CEF cell line.
Formulation:	State: Serum State: Lyophilized serum
Reconstitution Method:	Restore in distilled water.
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at -20°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	MAF bZIP transcription factor
Database Link:	<a href="#">Entrez Gene 4094 Human O75444</a>



[View online »](#)

**Background:**

c-maf is the cellular counterpart of oncogenic v-maf that belongs to the family of basic region leucine zipper domain transcription factors. The leucine-zipper domain is involved in the interaction with LRPICD. There are two forms of human c-maf mRNA, c-maf-long and c-maf-short. It is identified in the genome of the acute transforming avian retrovirus AS42. c-maf targets are IL-4 in Th2 cells, the crystalline genes in lens fiber cells, insulin gene in islet, p53 and L7 where it exerts its transcriptional role through binding to a Maf recognition element (MARE). It regulates Th2 differentiation and lineage-specific hematopoiesis. c-maf is a transcription factor for IL-10 gene expression in LPS-activated macrophages. Chromosomal aberration involving maf is found in some forms of multiple myeloma. It is expressed in myeloma cell lines and resting monocytes/macrophages.

**Synonyms:**

Transcription factor Maf, Proto-oncogene c-maf, V-maf

**Product images:**