

### **Product datasheet for AP55695PU-S**

#### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## beta Arrestin 1 (ARRB1) pSer412 Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1:500~1:1000.

Immunohistochemistry on paraffin sections: 1:50~1:100.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: Peptide sequence around phosphorylation site of Serine 412 (T-G-S(p)-P-Q) derived from

Human Arrestin 1 (KLH-conjugated)

**Specificity:** The antibody detects endogenous levels of Arrestin-1 only when phosphorylated at serine

412.

Formulation: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl,

0.02% sodium azide and 50% glycerol

State: Aff - Purified State: Liquid Ig fraction

**Concentration:** lot specific

**Purification:** Affinity chromatography using epitope-specific peptide

**Conjugation:** Unconjugated

Storage: Upon receipt, store undiluted (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 50 kDa

**Gene Name:** arrestin beta 1

Database Link: Entrez Gene 408 Human

P49407




#### beta Arrestin 1 (ARRB1) pSer412 Rabbit Polyclonal Antibody – AP55695PU-S

Background:


Functions in regulating agonist-mediated G-protein coupled receptor (GPCR) signaling by mediating both receptor desensitization and resensitization processes. During homologous desensitization, beta-arrestins bind to the GPRK-phosphorylated receptor and sterically preclude its coupling to the cognate G-protein; the binding appears to require additional receptor determinants exposed only in the active receptor conformation.

**Synonyms:** ARR1, Beta-arrestin-1

# **Product images:**



Western blot analysis of extracts from COS7 cells treated with Etoposide using Arrestin 1 (Phospho-Ser412) Antibody. The lane on the right is treated with the antigen-specific peptide.



Immunohistochemical analysis of paraffinembedded human breast carcinoma tissue using Arrestin 1 (Phospho-Ser412) antibody (left)or the same antibody preincubated with blocking peptide (right).