

Product datasheet for AP55709PU-N

PEA15 pSer104 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1:500~1:1000.

Immunohistochemistry on paraffin sections: 1:50~1:100.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Peptide sequence around phosphorylation site of Serine 104(I-P-S(p)-A-K) derived from

Human PEA-15 (KLH-conjugated)

Specificity: The antibody detects endogenous levels of PEA-15 only when phosphorylated at serine 104.

Formulation: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl,

0.02% sodium azide and 50% glycerol

State: Aff - Purified State: Liquid Ig fraction

Concentration: lot specific

Purification: Affinity chromatography using epitope-specific peptide

Conjugation: Unconjugated

Storage: Upon receipt, store undiluted (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 19 kDa

Gene Name: phosphoprotein enriched in astrocytes 15

Database Link: Entrez Gene 8682 Human

Q15121



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Blocks Ras-mediated inhibition of integrin activation and modulates the ERK MAP kinase cascade. Inhibits RPS6KA3 activities by retaining it in the cytoplasm. Inhibits both TNFRSF6-and TNFRSF1A-mediated CASP8 activity and apoptosis. Regulates glucose transport by controlling both the content of SLC2A1 glucose transporters on the plasma membrane and the insulin-dependent trafficking of SLC2A4 from the cell interior to the surface.


Synonyms:

Astrocytic phosphoprotein PEA-15, PED

Product images:



Western blot analysis of extracts from COS cells treated with TNF using PEA-15 (Phospho-Ser104) Antibody. The lane on the right is treated with the antigen-specific peptide.



Immunohistochemical analysis of paraffinembedded human heart tissue using PEA-15 (Phospho-Ser104) antibody (left)or the same antibody preincubated with blocking peptide (right).