

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for AP55784PU-S

## CD22 pTyr807 Rabbit Polyclonal Antibody

## **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western blot: 1:500~1:1000.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide sequence around phosphorylation site of tyrosine 807 (G-D-Y(p)-E-N) derived from Human BL-CAM (KLH-conjugated)
Specificity:	The antibody detects endogenous levels of BL-CAM only when phosphorylated at tyrosine 807.
Formulation:	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol State: Aff - Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Affinity chromatography using epitope-specific peptide
Conjugation:	Unconjugated
Storage:	Upon receipt, store undiluted (in aliquots) at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	95 kDa
Gene Name:	CD22 molecule
Database Link:	<u>Entrez Gene 933 Human</u> <u>P20273</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

# CD22 pTyr807 Rabbit Polyclonal Antibody - AP55784PU-SBackground:Mediates B-cell B-cell interactions. May be involved in the localization of B-cells in lymphoid<br/>tissues. Binds sialylated glycoproteins; one of which is CD45. Preferentially binds to alpha2,6-<br/>linked sialic acid. The sialic acid recognition site can be masked by cis interactions with sialic<br/>acids on the same cell surface. Upon ligand induced tyrosine phosphorylation in the immune<br/>response seems to be involved in regulation of B cell antigen receptor signaling. Plays a role<br/>in positive regulation through interaction with Src family tyrosine kinases and may also act as<br/>an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that<br/>block signal transduction through dephosphorylation of signaling molecules.

### Synonyms: SIGLEC2, Siglec-2, B-cell receptor CD22, Leu-14, BL-CAM

# **Product images:**



Western blot analysis of extracts from COS7 cells (Lane 2), using BL-CAM (Phospho-Tyr807) Antibody.The lane on the left is treated with antigen-specific peptide.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US