

## Product datasheet for **AP55800PU-S**

### AIRE pSer156 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	<b>Western blot:</b> 1:500~1:1000.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Peptide sequence around phosphorylation site of Serine156 P-G-S(p)-Q-L) derived from Human AIRE (KLH-conjugated)
Specificity:	The antibody detects endogenous levels of AIRE only when phosphorylated at serine 156.
Formulation:	Rabbit IgG in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol State: Aff - Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Affinity chromatography using epitope-specific peptide
Conjugation:	Unconjugated
Storage:	Upon receipt, store undiluted (in aliquots) at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	50 kDa
Gene Name:	autoimmune regulator
Database Link:	<a href="#">Entrez Gene 326 Human O43918</a>




[View online »](#)

**Background:**

The function of the protein encoded by this gene is not well defined, however it contains zinc finger motifs suggestive of a transcription factor. The protein (isoform 1) is localized to both the nucleus and cytoplasm. Three splice variant mRNAs products have been described [1]. The longer AIRE-1 mRNA appears to be more abundant and includes exons 1 through 14. Splice variant AIRE-2 includes a portion of the non-coding region of exon 1, an alternatively spliced longer exon 8, plus exons 9 through 14. Variant AIRE-3 includes the same exon 1-8-9 sequences as found in AIRE-2 but utilizes additional alternative splicing in exon 10 that shifts the reading frame such that a stop codon in exon 12 is utilized.

**Synonyms:**

Autoimmune regulator

**Product images:**


Western blot analysis of extracts from HeLa cells treated with Hu using AIRE (Phospho-Ser156) Antibody. The lane on the right is treated with the antigen-specific peptide.