

Product datasheet for AP55824PU-S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

MDC1 pSer513 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: Immunohistochemistry on paraffin sections: 1:50~1:100.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: Peptide sequence around phosphorylation site of Serine 513(E-R-S(p)-Q-A) derived from

Human MDC1 (KLH-conjugated)

Specificity: The antibody detects endogenous levels of MDC1 only when phosphorylated at serine 513.

Formulation: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl,

0.02% sodium azide and 50% glycerol

State: Aff - Purified State: Liquid Ig fraction

Concentration: lot specific

Purification: Affinity chromatography using epitope-specific peptide

Conjugation: Unconjugated

Storage: Upon receipt, store undiluted (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 226 kDa

Gene Name: mediator of DNA damage checkpoint 1

Database Link: Entrez Gene 9656 Human

Q14676

Background: The MDC1 protein contains a forkhead homology-associated (FHA) domain and 2 BRCA1

(MIM 113705) C-terminal motifs (BRCTs) and is required for the intra-S-phase DNA damage


checkpoint.


Synonyms: MDC1, KIAA0170, NFBD1





Product images:





Immunohistochemical analysis of paraffinembedded human brain tissue using MDC1 (Phospho-Ser513) Antibody (left)or the same antibody preincubated with blocking peptide (right).