

Product datasheet for AP55846PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

MCSF Receptor (CSF1R) pTyr708 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: Western blot: 1:500~1:1000.

Reactivity: Human, Mouse, Rat

Host: Rabbit
Clonality: Polyclonal

Immunogen: Peptide sequence around phosphorylation site of tyrosine 708 (K-K-Y(p)-V-R) derived from

Human CSFR (KLH-conjugated)

Specificity: The Antibody detects endogenous levels of CSF1R only when phosphorylated at tyrosine 708.

Formulation: Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl,

0.02% sodium azide and 50% glycerol

State: Aff - Purified State: Liquid Ig fraction

Concentration: lot specific

Purification: Affinity chromatography using epitope-specific peptide

Conjugation: Unconjugated

Storage: Upon receipt, store undiluted (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 200 kDa

Gene Name: colony stimulating factor 1 receptor

Database Link: Entrez Gene 1436 Human

P07333




Background:

Tyrosine-protein kinase that acts as cell-surface receptor for CSF1 and IL34 and plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes, such as macrophages and monocytes. Promotes the release of proinflammatory chemokines in response to IL34 and CSF1, and thereby plays an important role in innate immunity and in inflammatory processes. Plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone and tooth development.

Synonyms:

CSF-1-R, Fms proto-oncogene, c-fms, CSF1R, FMS, Macrophage colony-stimulating factor 1 receptor

Product images:



Western blot analysis of extracts from HepG2 cells using CSFR (Phospho-Tyr708) Antibody.The lane on the right is treated with the antigen-specific peptide.