

Product datasheet for **AP60004PU-L**

Egfl7 (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IP, WB
Recommended Dilution:	Western blot: 1-5 µg/ml. Immunocytochemistry. Immunoprecipitation.
Reactivity:	Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Highly pure recombinant mouse EGFL7 (Glu181-Leu275; C-terminus) derived from E. coli
Specificity:	This antibody recognizes VE-Statin/EGFL7.
Formulation:	PBS State: Purified State: Lyophilized purified IgG fraction
Reconstitution Method:	Restore in sterile water to a concentration of 0.1-1.0 mg/ml.
Purification:	Protein A chromatography
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	EGF-like domain 7
Database Link:	Entrez Gene 353156 Mouse Q9QXT5



[View online »](#)


Background:

EGFL7 (EGF-like domain-containing protein 7) regulates vascular tubulogenesis in vivo. During sprouting angiogenesis, groups of endothelial cells (ECs) migrate together in units called sprouts. EGFL7 regulates the proper spatial organization of ECs within each sprout and influences their collective movement. It inhibits platelet-derived growth factor (PDGF)-BB-induced smooth muscle cell migration and promotes endothelial cells adhesion to the substrate in vitro.


EGFL7 is an ~ 30 kDa secreted protein that contain an Emilin-like (EMI) domain (a multimerization motif), and 2 epidermal growth factor (EGF) domains, one of which binds calcium. Based on these domains, it has been hypothesized that EGFL7 may self-assemble like extracellular matrix (ECM) proteins and, thus, could incorporate into ECM. EGFL7 has been reported to stimulate cell adhesion as well as motility in a manner similar to ECM proteins. EGFL7 has been shown to be primarily expressed by developing ECs but also by primordial germ cells and some central nervous system neurons. Interestingly, EGFL7 expression markedly decreases in ECs in postnatal life, but can be strongly up-regulated after various tissue injuries that lead to increased angiogenic responses.

Synonyms:


EGF-like protein 7, EGF like protein 7, VE-statin, NOTCH4-like protein, Zneu1, MEGF7, UNQ187, PRO1449

Product images:


Western analysis of recombinant human and mouse EGFL7 using anti-mouse EGFL7 antibody. There is no cross reactivity visible with human EGFL-7. Performed by Dr. Frank Bicker, Rresearch group ??Molecular Signal Transduction?? (Prof. Dr. Mirko HH Schmidt), Institute of Microscopic Anatomy and Neurobiology, University Mainz, Germany.



Immunocytochemical staining (ICC) of human and mouse EGFL7: Left column: DAPI; Middle column: Staining with anti-mouse EGFL7 antibody.; Right column: Merge. Performed by Dr. Frank Bicker, Research group ??Molecular Signal Transduction?? (Prof. Dr. Mirko HH Schmidt), Institute of Microscopic Anatomy and Neurobiology, University Mainz, Germany.



Immunoprecipitation of human and mouse EGFL7 constructs with anti-mouse EGFL7 antibody. and subsequent Western analysis with anti-V5 antibodies. Samples were loaded in 15% SDS-polyacrylamide gel under reducing conditions. Performed by Dr. Frank Bicker, Rresearch group ??Molecular Signal Transduction?? (Prof. Dr. Mirko HH Schmidt), Institute of Microscopic Anatomy and Neurobiology, University Mainz, Germany.