

Product datasheet for **AP60005PU-L**

PRAME (C-term) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	Western blot: 1-5 µg/ml. ELISA: 0.1-1 µg/ml.
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Highly pure (>95%) recombinant human PAME /MAPE C-terminal end (Met321-Asn509) derived from <i>E. coli</i>
Specificity:	This antibody detects PRAME / MAPE at C-term.
Formulation:	PBS State: Purified State: Lyophilized purified Ig fraction
Reconstitution Method:	Centrifuge vial prior to opening. Restore in sterile water to a concentration of 0.1-1.0 mg/ml.
Purification:	Protein A chromatography
Conjugation:	Unconjugated
Storage:	Store lyophilized at 2-8°C for 6 month or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	preferentially expressed antigen in melanoma
Database Link:	Entrez Gene 23532 Human P78395




[View online »](#)

Background:


PRAME/MAPE/OIP4 is a germinal tissue-specific gene that is also expressed at high levels in haematological malignancies and solid tumors. The physiological functions of PRAME in normal and tumor cells are unknown, although a role in the regulation of retinoic acid signaling has been proposed. Sequence homology and structural predictions suggest that PRAME is related to the Leucine-rich repeat (LRR) family of proteins, which have diverse functions. PRAME, or „preferentially expressed antigen in melanoma”, was originally identified as a gene encoding a HLA-A24 restricted antigenic peptide presented to autologous tumor-specific cytotoxic T lymphocytes derived from a patient with melanoma. PRAME is synonymous with MAPE (melanoma antigen preferentially expressed in tumors) and OIP4 (OPA-interacting protein 4), and its expression profile defines it as a cancer-testis antigen. Cancer-testis antigens (CTAs) are encoded by non-mutated genes expressed at high levels in germinal tissues and tumors, but which are absent from or detected at low levels in other tissues. PRAME may be somewhat different to other cancer-testis antigens in that it shows some expression in normal tissues such as ovary, adrenal, placenta and endometrium. The C-terminus of human PRAME (amino acids 453-509) was also identified to bind *Neisseria gonorrhoeae* opacity factors, in this case the OPA-P protein. Thus PRAME is also known as OIP4 (OPA interacting protein).

Synonyms:

OPA-interacting protein 4, OIP4, OIP-4

Product images:

Western analysis using anti-human PRAME / MAPE antibody. Sample was loaded in 21% SDS-polyacrylamide gel under reducing conditions. Lane 1: MWM (kDa); Lane 2: rh PRAME.



ELISA using the polyclonal rabbit anti-human PRAME / M-N) as standard.