

Product datasheet for AR00147PU-N

Toxoplasma gondii p29 (GRA7) Apicomplexan Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Recombinant Proteins
Description:	Toxoplasma gondii p29 (GRA7) recombinant protein, 1 mg
Species:	Apicomplexan
Expression Host:	E. coli
Concentration:	lot specific
Purity:	>95% pure by 10% PAGE, coomassie staining. Purification: GS-4B Sepharose-Affinity Chromatography.
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 50 mM Tris-HCl, 1.5M Urea containing 50% Glycerol without preservatives
Preparation:	Liquid purified protein
Applications:	ELISA. Western Blot.
Protein Description:	Recombinant Protein Toxoplasma gondii P29 (GRA7). Contains the P29 (GRA7) immunodominant region. Contains GST fusion partner. Immunoreactive with Toxoplasma gondii positive sera.
Storage:	Store the protein at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>XP 002367613</u>
Synonyms:	Dense granule protein 7



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Toxoplasma gondii p29 (GRA7) Apicomplexan Protein – AR00147PU-N
Summary:	Toxoplasma is a crescent shaped sporozoan that lives as an intracellular parasite in various tissues of many vertebrates and completes its life cycle in a single host. Its life cycle includes two phases called the intestinal (or enteroepithelial) and extraintestinal phases. The intestinal phase produces oocysts and occurs only in cats, wild as well as domesticated. The extraintestinal phase occurs in all infected animals including cats, and produces tachyzoites (actively proliferating trophozoites) and eventually, bradyzoites (slowly growing trophozoites) or zoitocysts. Infection due to Toxoplasma gondii occurs in pregnant women where a variable degree of immunosuppression may exist or in patients receiving immunosuppressive drug therapy. Toxoplasma infects tissue of the GI tract where an active infection is accompanied by fever and enlargement of the spleen. Symptoms of toxoplasmosis, in which the maternal infection is transmitted during pregnancy, can produce blindness or mental retardation in the newborn.
Protein Families	s: ELISA. Western Blot.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US