

## Product datasheet for **AR03020PU-L**

### HSPA8 / HSC70 (active) Human Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	HSPA8 / HSC70 (active) human recombinant protein, 2x0.1 mg
Species:	Human
Expression Host:	E. coli
Concentration:	lot specific
Purity:	>90% pure as determined by SDS-PAGE analysis.
Buffer:	Presentation State: Aff - Purified State: Liquid affinity purified protein Buffer System: Na-Phosphate, pH 7.5 (20 mM), 150 mM NaCl, 10% Glycerol, 200 mM Imidazole.
Preparation:	Liquid affinity purified protein
Applications:	ATPase Assay, WB control, Binding Assays, ELISA reference standard. This protein has ATPase activity at the time of manufacture of 3.2 uM phosphate liberated/hr/ug protein in a 200 uL reaction at 37 C (pH 7.5) in the presence of 20 uL of 1 mM ATP using a Malachite Green assay.
Protein Description:	Recombinant Human Hsc70 Protein with ATPase activity, his-tagged
Storage:	Store the antibody (in aliquots) at -20°C. Can be shipped at 2-8 °C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
RefSeq:	<a href="#">NP_006588</a>
Locus ID:	3312
UniProt ID:	<a href="#">P11142</a>
Cytogenetics:	11q24.1
Synonyms:	HSP73, HSPA10, Heat shock cognate 71 kDa protein, Heat shock 70 kDa protein 8



[View online »](#)

- Summary:** This gene encodes a member of the heat shock protein 70 family, which contains both heat-inducible and constitutively expressed members. This protein belongs to the latter group, which are also referred to as heat-shock cognate proteins. It functions as a chaperone, and binds to nascent polypeptides to facilitate correct folding. It also functions as an ATPase in the disassembly of clathrin-coated vesicles during transport of membrane components through the cell. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2011]
- Protein Families:** ATPase Assay, WB control, Binding Assays, ELISA reference standard.  
This protein has ATPase activity at the time of manufacture of 3.2 uM phosphate liberated/hr/ug protein in a 200 uL reaction at 37 C (pH 7.5) in the presence of 20 uL of 1 mM ATP using a Malachite Green assay.
- Protein Pathways:** Antigen processing and presentation, Endocytosis, MAPK signaling pathway, Spliceosome