

Product datasheet for **AR08013TR-N**

Bovine IgG (Fc Fragment) Bovine Protein

Product data:

Product Type:	Native Proteins
Description:	Bovine IgG Fc Fragment bovine protein, 1 mg
Species:	Bovine
Protein Source:	Serum
Concentration:	lot specific
Purity:	>95% normal serum delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above
Conjugation:	Texas Red
Buffer:	State: Lyophilized IgG F(c) fragment Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Polyethylene Glycol and 0.01% (w/v) Sodium Azide <u>Label:</u> Texas Red(TM) Sulfonyl Chloride (Molecular Weight 625 daltons) <u>Absorption/Emission:</u> 596 nm / 620 nm <u>Protein Ratio:</u> 3.1 moles Texas Red(TM) per mole of Bovine IgG F(c) Presentation Label: Texas Red
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent). For extended storage add glycerol to 50%
Preparation:	Lyophilized IgG F(c) fragment
Applications:	Immunofluorescence. Flow Cytometry.
Protein Description:	Texas Red(TM) Conjugated Bovine IgG F(c) fragment.
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store the protein undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.



[View online »](#)