

Product datasheet for AR09000PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

Adiponectin (18-247) Mouse Protein

Product data:

Product Type: Recombinant Proteins

Description: Adiponectin (18-247) mouse recombinant protein, 0.5 mg

Species: Mouse Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MGSSHHHHHH SSGLVPRGSH MEDDVTTTEE LAPALVPPPK GTCAGWMAGI PGHPGHNGTP

GRDGRDGTPG EKGEKGDAGL LGPKGETGDV GMTGAEGPRG FPGTPGRKGE PGEAAYVYRS
AFSVGLETRV TVPNVPIRFT KIFYNQQNHY DGSTGKFYCN IPGLYYFSYH ITVYMKDVKV SLFKKDKAVL

FTYDQYQEKN VDQASGSVLL HLEVGDQVWL QVYGDGDHNG LYADNVNDST FTGFLLYHDT N

Predicted MW: 27 kDa

Concentration: lot specific

Purity: >90% > / = 90% by SDS-PAGE

Buffer: Presentation State: Purified

State: Liquid

Buffer System: 20 mM Tris pH 8.0, 1 mM DTT, 10% Glycerol

Preparation: Liquid

Protein Description: Recombinant mouse adiponectin, fused to His-tag at N-terminus, was expressed in E.coli and

purified by using conventional chromatography techniques.

Storage: Store at -20°C. Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 033735

 Locus ID:
 11450

 UniProt ID:
 Q60994

Cytogenetics: 16 13.96 cM

Synonyms: 30kDa; Acdc; Acrp30; Ad; adipo; apM1; APN; GBP28





Summary:

Important adipokine involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. Inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. May play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex, LMW, MMW or HMW.[UniProtKB/Swiss-Prot Function]

Product images:

