

Product datasheet for **AR09015PU-N**

Cyclophilin A (1-165, His-tag) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Cyclophilin A (1-165, His-tag) human recombinant protein, 0.1 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	<u>MGSSHHHHHH SSGLVPRGSH</u> MVNPTVFFDI AVDGEPLGRV SFELFADKVP KTAENFRALS TGEKGFQYKQ SCFHRIIPGF MCQGGDFTRH NGTGGKSIYG EKFDENFIL KHTGPGILSM ANAGPNTNGS QFFICTAKTE WLDGKHVVFG KVKEGMNIVE AMERFGSRNG KTSKKITIAD CGQLE
Tag:	His-tag
Predicted MW:	20 kDa
Concentration:	lot specific
Purity:	>95% by SDS PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl (pH 8.0) containing 20 mM NaCl, 0.5 mM DTT, 10% glycerol
Bioactivity:	Specific: > 270 nmoles/min/mg, defined as the amount of enzyme that cleaves 1 umole of suc-AAPF-pNA per minute at 25°C in Tris-HCl pH 8.0 using chymotrypsin.
Endotoxin:	< 1.0 EU per 1 µg of protein (determined by LAL method)
Preparation:	Liquid purified protein
Applications:	Protocol: Activity Assay 1. Prepare 170 ul assay buffer into a suitable container and pre-chill on ice before use: The final concentrations are 200 mM Tris-HCl, pH 8.0, and 20nM chymotrypsin. 2. Add 10 ul of recombinant Cyclophilin A protein with 1 ug in assay buffer. 3. Mix by inversion and equilibrate to 1°C and monitor the A405nm until the value is constant using a spectrophotometer. 4. Add 20 ul pre-chilled 5mM suc-AAFP-pNA. (Substrate was dissolved in TFE that contained 460mM LiCl to a concentration of 3 mM) 5. Record the increase in A405 nm for 30 minutes at 25°C.
Protein Description:	Recombinant Human PPIA, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.



[View online »](#)

Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	NP_001287910
Locus ID:	5478
UniProt ID:	P62937 , Q567Q0
Cytogenetics:	7p13
Synonyms:	CYPA; CYPH; HEL-S-69p
Summary:	This gene encodes a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. The encoded protein is a cyclosporin binding-protein and may play a role in cyclosporin A-mediated immunosuppression. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions. Multiple pseudogenes that map to different chromosomes have been reported. [provided by RefSeq, Jul 2008]

Product images:

