

Product datasheet for AR09017PU-L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Desert hedgehog / DHH (23-198, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: Desert hedgehog / DHH (23-198, His-tag) human recombinant protein, 0.5 mg

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MGSSHHHHHH SSGLVPRGSH MCGPGRGPVG RRRYARKQLV PLLYKQFVPG VPERTLGASG PAEGRVARGS ERFRDLVPNY NPDIIFKDEE NSGADRLMTE RCKERVNALA IAVMNMWPGV RLRVTEGWDE DGHHAODSLH YEGRALDITT SDRDRNKYGL LARLAVEAGF DWVYYESRNH

VHVSVKADNS LAVRAGG

Tag: His-tag
Predicted MW: 22 kDa

Concentration: lot specific

Purity: >95% > 95% by SDS PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM MES pH 5.5 containing 0.5 mM DTT, 20% Glycerol

Preparation: Liquid purified protein

Protein Description: Recombinant DHH protein, fused to His-tag, was expressed in E.coli and purified by using

conventional chromatography techniques.

Note: NCBI Accession No.: NP_066382

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 066382

 Locus ID:
 50846

 UniProt ID:
 043323

 Cytogenetics:
 12q13.12

Synonyms: GDMN; GDXYM; HHG-3; SRXY7





Summary:

This gene encodes a member of the hedgehog family. The hedgehog gene family encodes signaling molecules that play an important role in regulating morphogenesis. This protein is predicted to be made as a precursor that is autocatalytically cleaved; the N-terminal portion is soluble and contains the signalling activity while the C-terminal portion is involved in precursor processing. More importantly, the C-terminal product covalently attaches a cholesterol moiety to the N-terminal product, restricting the N-terminal product to the cell surface and preventing it from freely diffusing throughout the organism. Defects in this protein have been associated with partial gonadal dysgenesis (PGD) accompanied by minifascicular polyneuropathy. This protein may be involved in both male gonadal differentiation and perineurial development. [provided by RefSeq, May 2010]

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protease

Protein Pathways: Hedgehog signaling pathway

Product images:

