

Product datasheet for **AR09072PU-L**

Cyclophilin H / PPIH (1-177) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Cyclophilin H / PPIH (1-177) human recombinant protein, 0.5 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MAVANSSPVN PVVFFDVSIG GQEVGRMKIE LFADVVPKTA ENFRQFCTGE FRKDGVPIGY KGSTFHRVIK DFMIQGGDFV NGDGTGVASI YRGPFADEFN KLRHSAPGLL SMANSGPSTN GCQFFITCSK CDWLDGKHVV FGKIIDGLLV MRKIENVPTG PNNKPKLPVV ISQCGEM
Predicted MW:	19.2 kDa
Concentration:	lot specific
Purity:	>95% by SDS PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: PBS, pH 7.4, 10% glycerol
Bioactivity:	Specific: > 220 nmoles/min/mg, defined as the amount of enzyme that cleaves 1 umole of suc-AAPF-pNA per minute at 25°C in Tris-HCl pH 8.0 using chymotrypsin.
Endotoxin:	< 1.0 EU per 1 µg of protein (determined by LAL method)
Preparation:	Liquid purified protein
Applications:	Protocol: Activity Assay 1. Prepare 170 ul assay buffer into a suitable container and pre-chill on ice before use: The final concentrations are 200 mM Tris-HCl, pH 8.0, and 20nM chymotrypsin. 2. Add 10 ul of recombinant Cyclophilin H (PPIH) protein with 1 ug in assay buffer. 3. Mix by inversion and equilibrate to 1°C and monitor the A405nm until the value is constant using a spectrophotometer. 4. Add 20 ul pre-chilled 5mM suc-AAFP-pNA. (Substrate was dissolved in TFE that contained 460mM LiCl to a concentration of 3 mM) 5. Record the increase in A405 nm for 30 minutes at 25°C.
Protein Description:	Recombinant human cyclophilin H was expressed in E.coli and purified by conventional chromatography techniques.
Storage:	Store (in aliquots) at -20°C. Avoid repeated freezing and thawing.



[View online »](#)

Stability: Shelf life: one year from despatch.

RefSeq: [NP_001317439](#)

Locus ID: 10465

UniProt ID: [O43447](#)

Cytogenetics: 1p34.2

Synonyms: CYP-20; CYPH; SnuCyp-20; USA-CYP

Summary: The protein encoded by this gene is a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. This protein is a specific component of the complex that includes pre-mRNA processing factors PRPF3, PRPF4, and PRPF18, as well as U4/U5/U6 tri-snRNP. This protein has been shown to possess PPIase activity and may act as a protein chaperone that mediates the interactions between different proteins inside the spliceosome. [provided by RefSeq, Jul 2008]

Protein Pathways: Spliceosome

Product images:

