

Product datasheet for AR09089PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Thioredoxin / TRX1 (1-105) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: Thioredoxin / TRX1 (1-105) human recombinant protein, 0.1 mg

Species: Human
Expression Host: E. coli

Expression cDNA Clone MVKQIESKTA FQEALDAAGD KLVVVDFSAT WCGPCKMIKP FFHSLSEKYS NVIFLEVDVD

or AA Sequence: DCQDVASECE VKCMPTFQFF KKGQKVGEFS GANKEKLEAT INELV

Predicted MW: 11.7 kDa

Concentration: lot specific

Purity: ≥90% by SDS PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein Buffer System: PBS, pH 7.4

Bioactivity: Specific activity is >150 A650/cm/min/mg, obtained by measuring the increase of insulin

precipitation in absorbance at 650 nm resulting from the reduction of insulin. Please refer to

our activity assay protocol below.

Preparation: Liquid purified protein

Applications: Protocol: Activity Assay

1. Prepare a 1 ml reaction cocktail into a suitable container: The final concentrations are

5mM Tris, 63mM sodium phosphate, 1mM DTT, 0.1% (w/v) insulin, 2mM EDTA. 2. Add 4ug of recombinant thioredoxin into reaction cocktail and mix immediately.

3. Incubate at 25°C for 20 minutes.

4. Record the increase in A650nm for 15 minutes

Protein Description: Recombinant human thioredoxin-1 was overexpressed in E. coli and purified by using

conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001231867





 Locus ID:
 7295

 UniProt ID:
 P10599

 Cytogenetics:
 9q31.3

Synonyms: TRDX; TRX; TRX1; Trx80

Summary: The protein encoded by this gene acts as a homodimer and is involved in many redox

reactions. The encoded protein is active in the reversible S-nitrosylation of cysteines in certain proteins, which is part of the response to intracellular nitric oxide. This protein is found in the cytoplasm. Two transcript variants encoding different isoforms have been found

for this gene. [provided by RefSeq, Oct 2011]

Protein Families: Druggable Genome

Product images:

