

### **Product datasheet for AR09149PU-N**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## EIF5A1 (1-154) Human Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** EIF5A1 (1-154) human recombinant protein, 0.1 mg

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MADDLDFETG DAGASATFPM QCSALRKNGF VVLKGRPCKI VEMSTSKTGK HGHAKVHLVG IDIFTGKKYE DICPSTHNMD VPNIKRNDFQ LIGIQDGYLS LLQDSGEVRE DLRLPEGDLG

KEIEQKYDCG EEILITVLSA MTEEAAVAIK AMAK

Predicted MW: 16.8 kDa

Concentration: lot specific

Purity: >95% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 50 mM Tris-HCl buffer pH 7.5 containing 10% glycerol

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant human eIF5A protein was expressed in E. coli and purified by using

conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**RefSeq:** NP 001137232

**Locus ID:** 1984

UniProt ID: <u>P63241</u>, <u>P63241-2</u>

Cytogenetics: 17p13.1

**Synonyms:** eIF-4D; EIF-5A; EIF5A1; eIF5Al





#### **Summary:**

mRNA-binding protein involved in translation elongation. Has an important function at the level of mRNA turnover, probably acting downstream of decapping. Involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall integrity. With syntenin SDCBP, functions as a regulator of p53/TP53 and p53/TP53-dependent apoptosis. Regulates also TNF-alphamediated apoptosis. Mediates effects of polyamines on neuronal process extension and survival. May play an important role in brain development and function, and in skeletal muscle stem cell differentiation. Also described as a cellular cofactor of human T-cell leukemia virus type I (HTLV-1) Rex protein and of human immunodeficiency virus type 1 (HIV-1) Rev protein, essential for mRNA export of retroviral transcripts.[UniProtKB/Swiss-Prot Function]

#### **Product images:**

