

## Product datasheet for **AR09190PU-L**

### CHGA (Vasostatin-2 chain) (19-131) Human Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	CHGA (Vasostatin-2 chain) (19-131) human recombinant protein, 0.5 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MLPVNSPMNK GDTEVMKCIV EVISDTLSKP SPMPVSQECF ETLRGDERIL SILRHQNLK ELQDLALQGA KERAHQQKH SGFEDELSEV LENQSSQAEK KEAVEEPSSK DVME
Predicted MW:	12.8 kDa
Concentration:	lot specific
Purity:	>95% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 10% glycerol
Endotoxin:	< 1.0 EU per 1 µg of protein (determined by LAL method )
Preparation:	Liquid purified protein
Protein Description:	Recombinant human Vasostatin 2 protein was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<a href="#">NP_001266</a>
Locus ID:	1113
UniProt ID:	<a href="#">P10645</a> , <a href="#">Q86T07</a>
Cytogenetics:	14q32.12
Synonyms:	CGA



[View online »](#)

**Summary:**

The protein encoded by this gene is a member of the chromogranin/secretogranin family of neuroendocrine secretory proteins. It is found in secretory vesicles of neurons and endocrine cells. This gene product is a precursor to three biologically active peptides; vasostatin, pancreastatin, and parastatin. These peptides act as autocrine or paracrine negative modulators of the neuroendocrine system. Two other peptides, catestatin and chromofungin, have antimicrobial activity and antifungal activity, respectively. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2014]

**Protein Families:**

Druggable Genome, Secreted Protein

**Product images:**