

Product datasheet for **AR09205PU-N**

BMF (1-129, T7-tag) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	BMF (1-129, T7-tag) human recombinant protein, 0.1 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	<u>MASMTGGQOM</u> <u>GRGSHMEPSQ</u> CVEELEDVVF QPEDGEPVTQ PGSLLSADLF AQSLDCPLS RLQLFPLTHC CGPGLRPTSQ EDKATQTLSP ASPSQGVMLP CGVTEEPQRL FYAPAEPKSC VVADPPLPAQ PCFEWRREQE RGRP
Tag:	T7-tag
Predicted MW:	15.6 kDa
Concentration:	lot specific
Purity:	>90% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 7.5) containing, 1 mM DTT, 10% glycerol
Endotoxin:	< 1.0 EU per 1 µg of protein (determined by LAL method)
Preparation:	Liquid purified protein
Protein Description:	Recombinant human BMF, fused to T7-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography.
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>NP_001003940</u>
Locus ID:	90427
UniProt ID:	<u>Q96LC9</u>
Cytogenetics:	15q15.1



[View online »](#)

Summary:

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein contains a single BCL2 homology domain 3 (BH3), and has been shown to bind BCL2 proteins and function as an apoptotic activator. This protein is found to be sequestered to myosin V motors by its association with dynein light chain 2, which may be important for sensing intracellular damage and triggering apoptosis. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]

Protein Families:

Druggable Genome

Product images: