

## **Product datasheet for AR09261PU-N**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## Eotaxin-3 / CCL26 (24-94) Human Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Eotaxin-3 / CCL26 (24-94) human protein, 0.1 mg

Species: Human
Expression Host: E. coli

Expression cDNA Clone MTRGSDISKT CCFQYSHKPL PWTWVRSYEF TSNSCSQRAV IFTTKRGKKV CTHPRKKWVQ

or AA Sequence: KYISLLKTPK QL

Predicted MW: 8.5 kDa

Concentration: lot specific

**Purity:** >90% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified peptide

Buffer System: PBS, pH 7.4, containing 10% glycerol

**Endotoxin:** < 1.0 EU per 1 μg of protein (determined by LAL method )

**Preparation:** Liquid purified peptide

**Protein Description:** CCL26 was expressed in E.coli and purified by conventional chromatography, after refolding

of the isolated inclusion bodies in a renaturation buffer.

**Note:** (Real molecular weight on SDS-PAGE will be shift up).

**Storage:** Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

RefSeg: NP 006063

 Locus ID:
 10344

 UniProt ID:
 Q9Y258

 Cytogenetics:
 7q11.23

Synonyms: C-C motif chemokine 26, Small-inducible cytokine A26, SCYA26, MIP4 alpha, IMAC, TSC-1, TSC1





**Summary:** 

This gene is one of two Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 7. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for normal peripheral blood eosinophils and basophils. This protein also has antimicrobial activity, displaying an antibacterial effect on S. pneumoniae, S. aureus, Non-typeable H. influenzae, and P. aeruginosa. The product of this gene is one of three related chemokines that specifically activate chemokine receptor CCR3. This chemokine may contribute to the eosinophil accumulation in atopic diseases. [provided by RefSeq, Jul 2020]

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

## **Product images:**

