

Product datasheet for **AR09323PU-L**

BAG3 (1-575, His-tag) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	BAG3 (1-575, His-tag) human recombinant protein, 0.25 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	<u>MGSSHHHHHH SSGLVPRGSH</u> MSAATHSPMM QVASGNNGDRD PLPPGWEIKI DPQTGWPFV DHNSRTTTWN DPRVPSEGPK ETPSSANGPS REGSRLPPAR EGHPVYPQLR PGYIPIVLH EGAENRQVHP FHVYPQGMQ RFRTEAAAAA PQRSQSPLRG MPETTQPKDQ CGQVAAAAAA QPPASHGPER SQSPAASDCS SSSSSASLPS SGRSSLGSHQ LPRGYISIPV IHEQNVTRPA AQPSFHQAQK THYPAQQGEY QTHQPVYHKI QGDDWEPRPL RAASPFSSV QGASSREGSP ARSSTPLHSP SPIRVHTVVD RPQQPMTHRE TAPVSQPENK PESKPGVGP ELPPGHIPIQ VIRKEVDSKP VSQKPPPSE KVEVKVPPAP VPCPPSPGP SAVPSSPKSV ATEERAAPST APAEATPPPK GEAEAPPKHP GVLKVEAILE KVQGLEQAVD NFEKGKTDKK YLMIEEYLTK ELLALDSVDP EGRADVQRAR RDGVRKVQTI LEKLEQKAID VPGQVQVYEL QPSNLEADQP LQAIMEMGAV AADKGKKNAG NAEDPHTETQ QPEATAAATS NPSSMTDTPG NPAAP
Tag:	His-tag
Concentration:	lot specific
Purity:	>90% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 1 mM EDTA, 0.1 mM PMSF, 10% glycerol
Preparation:	Liquid purified protein
Protein Description:	Recombinant human BAG3, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>NP_004272</u>
Locus ID:	9531



[View online »](#)

UniProt ID: [O95817](#)

Cytogenetics: 10q26.11

Synonyms: BAG-3; BIS; CAIR-1; MFM6

Summary: BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome

Product images:

