

Product datasheet for **AR09348PU-N**

PDIA3 (25-505, His-tag) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	PDIA3 (25-505, His-tag) human recombinant protein, 0.1 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	<u>MRGSHHHHHH</u> <u>GMASMTGGQQ</u> <u>MGRDLYDDDD</u> <u>KDRWGMS</u> SDV LELTDDNFES RISDTGSAGL MLVEFFAPWC GHCKRLAPEY EAAATRLKGI VPLAKVDCTA NTNTCNKYGV SGYPTLKIFR DGEEAGAYDG PRTADGIVSH LKKQAGPASV PLRTEEFKK FISKDASIV GFFDDSFSEA HSEFLKAASN LRDNYRFAHT NVESLVNEYD DNGEGILFR PSHLTNKFED KTVAYTEQKM TSGKIKKFIQ ENIFGICPHM TEDNKDLIQG KDLLIAYYDV DYEKNAKGSN YWRNRVMMVA KKFLDAGHKL NFAVASRKTf SHELSDFGLE STAGEIPVVA IRTAKGEKfV MQEEFSRDGK ALERFLQDYF DGNLKRYLKS EPIPESNDGP VKVVAENFD EIVNNENKDV LIEFYAPWCG HCKNLEPKYK ELGEKLSKDP NIVIAKMDAT ANDVPSPYEV RGFPTIYFSP ANKKLNPKKY EGGRELSDFI SYLQREATNP PVIQEEKPKK KKKAQEDL
Tag:	His-tag
Predicted MW:	58.5 kDa
Concentration:	lot specific
Purity:	>95% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 1 mM DTT, 0.1 M NaCl, and 10% glycerol
Preparation:	Liquid purified protein
Protein Description:	Recombinant Human PDIA3 protein, fused to <i>His-tag</i> at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>NP_005304</u>
Locus ID:	2923



[View online »](#)

UniProt ID:	P30101 , V9HVV3
Cytogenetics:	15q15.3
Synonyms:	ER60; ERp57; ERp60; ERp61; GRP57; GRP58; HEL-S-93n; HEL-S-269; HsT17083; P58; PI-PLC
Summary:	This gene encodes a protein of the endoplasmic reticulum that interacts with lectin chaperones calreticulin and calnexin to modulate folding of newly synthesized glycoproteins. The protein was once thought to be a phospholipase; however, it has been demonstrated that the protein actually has protein disulfide isomerase activity. It is thought that complexes of lectins and this protein mediate protein folding by promoting formation of disulfide bonds in their glycoprotein substrates. This protein also functions as a molecular chaperone that prevents the formation of protein aggregates. [provided by RefSeq, Dec 2016]
Protein Families:	Druggable Genome
Protein Pathways:	Antigen processing and presentation

Product images: