

## **Product datasheet for AR09477PU-N**

## OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## DBI / ACBP (1-87, His-tag) Human Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** DBI / ACBP (1-87, His-tag) human recombinant protein, 0.1 mg

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** MGSSHHHHHH SSGLVPRGSH MSQAEFEKAA EEVRHLKTKP SDEEMLFIYG HYKQATVGDI

or AA Sequence: NTERPGMLDF TGKAKWDAWN ELKGTSKEDA MKAYINKVEE LKKKYGI

Tag: His-tag
Predicted MW: 12.2 kDa
Concentration: lot specific

**Purity:** >95% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.1 M NaCl, 10% glycerol

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant human ACBP protein, fused to His-tag at N-terminus, was expressed in E.coli

and purified by using conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**RefSeq:** NP 001073331

**Locus ID:** 1622

UniProt ID: <u>P07108</u>, <u>B8ZWD2</u>

**Cytogenetics:** 2q14.2

Synonyms: ACBD1; ACBP; CCK-RP; EP





**Summary:** 

This gene encodes diazepam binding inhibitor, a protein that is regulated by hormones and is involved in lipid metabolism and the displacement of beta-carbolines and benzodiazepines, which modulate signal transduction at type A gamma-aminobutyric acid receptors located in brain synapses. The protein is conserved from yeast to mammals, with the most highly conserved domain consisting of seven contiguous residues that constitute the hydrophobic binding site for medium- and long-chain acyl-Coenzyme A esters. Diazepam binding inhibitor is also known to mediate the feedback regulation of pancreatic secretion and the postprandial release of cholecystokinin, in addition to its role as a mediator in corticotropin-dependent adrenal steroidogenesis. Three pseudogenes located on chromosomes 6, 8 and 16 have been identified. Multiple transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome

**Protein Pathways:** PPAR signaling pathway

## **Product images:**

