

# Product datasheet for AR09556PU-N

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Alpha-Synuclein / SNCA (1-140) Mouse Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Alpha-Synuclein / SNCA (1-140) mouse recombinant protein, 0.1 mg

Species: Mouse Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

Clone MDVFMKGLSK AKEGVVAAAE KTKQGVAEAA GKTKEGVLYV GSKTKEGVVH GVTTVAEKTK EQVTNVGGAV VTGVTAVAQK TVEGAGNIAA ATGFVKKDQM GKGEEGYPQE GILEDMPVDP

GSEAYEMPSE EGYQDYEPEA

Predicted MW: 14.4 kDa

Concentration: lot specific

Purity: >95% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 7.5) containing 10% glycerol

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant mouse alpha-Synuclein was expressed in E.coli and purified by using

conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**RefSeq:** NP 001035916

 Locus ID:
 20617

 UniProt ID:
 055042

 Cytogenetics:
 6 29.15 cM

 Synonyms:
 NACP, PARK1





#### **Summary:**

Neuronal protein that plays several roles in synaptic activity such as regulation of synaptic vesicle trafficking and subsequent neurotransmitter release. Participates as a monomer in synaptic vesicle exocytosis by enhancing vesicle priming, fusion and dilation of exocytotic fusion pores. Mechanistically, acts by increasing local Ca(2+) release from microdomains which is essential for the enhancement of ATP-induced exocytosis. Acts also as a molecular chaperone in its multimeric membrane-bound state, assisting in the folding of synaptic fusion components called SNAREs (Soluble NSF Attachment Protein REceptors) at presynaptic plasma membrane in conjunction with cysteine string protein-alpha/DNAJC5 (PubMed:20798282, PubMed:25246573). This chaperone activity is important to sustain normal SNARE-complex assembly during aging. Plays also a role in the regulation of the dopamine neurotransmission by associating with the dopamine transporter (DAT1) and thereby modulating its activity (By similarity).[UniProtKB/Swiss-Prot Function]

### **Product images:**

