

Product datasheet for AR09607PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DR1 / NC2-beta (1-176, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: DR1 / NC2-beta (1-176, His-tag) human recombinant protein, 0.1 mg

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MGSSHHHHHH SSGLVPRGSH MASSSGNDDD LTIPRAAINK MIKETLPNVR VANDARELVV NCCTEFIHLI SSEANEICNK SEKKTISPEH VIQALESLGF GSYISEVKEV LQECKTVALK RRKASSRLEN

LGIPEEELLR QQQELFAKAR QQQAELAQQE WLQMQQAAQQ AQLAAASASA SNQAGSSQDE

EDDDDI

Tag: His-tag
Predicted MW: 21.6 kDa
Concentration: lot specific

Purity: >85% by SDS-PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl Buffer (pH 8.0) containing 100 mM NaCl, 0.1 mM PMSF, 10%

Glycerol

Preparation: Liquid purified protein

Protein Description: Recombinant human DR1 protein, fused to His-tag at N-terminus, was expressed in E.coli and

purified by using conventional chromatography.

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001929

Locus ID: 1810

UniProt ID: <u>Q01658</u>, <u>Q658N3</u>

Cytogenetics: 1p22.1

Synonyms: NC2; NC2-BETA; NC2B; NCB2





Summary:

This gene encodes a TBP- (TATA box-binding protein) associated phosphoprotein that represses both basal and activated levels of transcription. The encoded protein is phosphorylated in vivo and this phosphorylation affects its interaction with TBP. This protein contains a histone fold motif at the amino terminus, a TBP-binding domain, and a glutamine-and alanine-rich region. The binding of DR1 repressor complexes to TBP-promoter complexes may establish a mechanism in which an altered DNA conformation, together with the formation of higher order complexes, inhibits the assembly of the preinitiation complex and controls the rate of RNA polymerase II transcription. [provided by RefSeq, Jul 2008]

Protein Families:

Transcription Factors

Product images:

