

Product datasheet for **AR09656PU-N**

BAT1 / UAP56 (1-428, His-tag) Human Protein

Product data:

| | |
|---------------------------------------|---|
| Product Type: | Recombinant Proteins |
| Description: | BAT1 / UAP56 (1-428, His-tag) human recombinant protein, 50 µg |
| Species: | Human |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | <u>MGSSHHHHHH SSGLVPRGSH</u> MAENDVDNEL LDYEDDEVET AAGGDGAEAP AKKDVKGSYV SIHSSGFRDF LLKPELLRAI VDCGFEPHSE VQHECIPQAI LGMDVLCQAK SGMGKTAVFV LATLQQLPEPV TGQVSVLVMC HTRELAFAQIS KEYERFSKYM PNVKVAVFFG GLSIKKDEEV LKKNCPHIVV GTPGRILALA RNKSLNLKHI KHFILDECDK MLEQLDMRRD VQEIFRMTPH EKQVMMFSAT LSKEIRPVCR KFMQDPMEIF VDETKLTLH GLQQYYVKLK DNEKNRKLFD LLDVLEFNQV VIFVKSQRC IALAQLLVEQ NFPAIAIHRG MPQEERLSRY QQFKDFQRRR LVATNLFGRG MDIERVNIAF NYDMPEDSDT YLHRVARAGR FGTKGLAITF VSDENDAKIL NDVQDRFEVN ISELPDEIDI SSYIEQTR |
| Tag: | His-tag |
| Predicted MW: | 51.1 kDa |
| Concentration: | lot specific |
| Purity: | >90% by SDS - PAGE |
| Buffer: | Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 20% glycerol, 0.1M NaCl, 1 mM DTT |
| Preparation: | Liquid purified protein |
| Protein Description: | Recombinant human BAT1 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques. |
| Storage: | Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| RefSeq: | <u>NP_004631</u> |
| Locus ID: | 7919 |
| UniProt ID: | <u>Q13838</u> , <u>A0A024RCM3</u> |



[View online »](#)

Cytogenetics: 6p21.33

Synonyms: BAT1; D6S81E; UAP56

Summary: This gene encodes a member of the DEAD box family of RNA-dependent ATPases that mediate ATP hydrolysis during pre-mRNA splicing. The encoded protein is an essential splicing factor required for association of U2 small nuclear ribonucleoprotein with pre-mRNA, and it also plays an important role in mRNA export from the nucleus to the cytoplasm. This gene belongs to a cluster of genes localized in the vicinity of the genes encoding tumor necrosis factor alpha and tumor necrosis factor beta. These genes are all within the human major histocompatibility complex class III region. Mutations in this gene may be associated with rheumatoid arthritis. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on both chromosomes 6 and 11. Read-through transcription also occurs between this gene and the upstream ATP6V1G2 (ATPase, H⁺ transporting, lysosomal 13kDa, V1 subunit G2) gene. [provided by RefSeq, Feb 2011]

Protein Pathways: Spliceosome

Product images:

