

# **Product datasheet for AR09677PU-N**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Cytidine deaminase (1-146, His-tag) Human Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Cytidine deaminase (1-146, His-tag) human recombinant protein, 0.1 mg

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MGSSHHHHHH SSGLVPRGSH MAQKRPACTL KPECVQQLLV CSQEAKQSAY CPYSHFPVGA

ALLTQEGRIF KGCNIENACY PLGICAERTA IQKAVSEGYK DFRAIAIASD MQDDFISPCG

ACRQVMREFG TNWPVYMTKP DGTYIVMTVQ ELLPSSFGPE DLQKTQ

Tag: His-tag

Predicted MW: 18.3 kDa

Concentration: lot specific

Purity: >90%

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl Buffer (pH 8.0) containing 1 mM DTT, 2 mM EDTA, 100 mM

NaCl, 40% Glycerol

**Bioactivity:** Specific: Specific activity is > 10,000 pmol/min/ug, and is defined as the amount of required to

deaminate 1.0 pmole of cytidine per min at pH 7.5 at 25°C

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant human CDA protein, fused to His-tag at N-terminus, was expressed in E.coli and

purified by using conventional chromatography.

**Storage:** Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**RefSeq:** <u>NP 001776</u>

Locus ID: 978

 UniProt ID:
 P32320

 Cytogenetics:
 1p36.12





#### Cytidine deaminase (1-146, His-tag) Human Protein - AR09677PU-N

Synonyms: CDA, CDD

Summary: This gene encodes an enzyme involved in pyrimidine salvaging. The encoded protein forms a


homotetramer that catalyzes the irreversible hydrolytic deamination of cytidine and deoxycytidine to uridine and deoxyuridine, respectively. It is one of several deaminases responsible for maintaining the cellular pyrimidine pool. Mutations in this gene are associated with decreased sensitivity to the cytosine nucleoside analogue cytosine arabinoside used in the treatment of certain childhood leukemias. [provided by RefSeq, Jul

2008]

**Protein Families:** Stem cell - Pluripotency

**Protein Pathways:** Drug metabolism - other enzymes, Metabolic pathways, Pyrimidine metabolism

## **Product images:**



15% SDS-PAGE (3ug)