

Product datasheet for **AR09749PU-L**

Syntenin-1 / SDCBP (1-298, His-tag) Human Protein

Product data:

| | |
|---------------------------------------|---|
| Product Type: | Recombinant Proteins |
| Description: | Syntenin-1 / SDCBP (1-298, His-tag) human recombinant protein, 0.5 mg |
| Species: | Human |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | <u>MGSSHHHHHH SSGLVPRGSH</u> MSLYPSLEDL KVDKVIQAQT AFSANPANPA ILSEASAPIP HDGNLYPRLY PELSQYMGLS LNEEEIRANV AVVSGAPLQG QLVARPSSIN YMVAVPTGND VGIRRAEIKQ GIREVILCKD QDGKIGLRLK SIDNGIFVQL VQANSPASLV GLRFGDQVLQ INGENCAGWS SDKAHKVLKQ AFGEKITMTI RDRPFERTIT MHKDSTGHVG FIFKNGKITS IVKDSSAARN GLLTEHNICE INGQNVIGLK DSQIADILST SGTWVTITIM PAFIFEHIK RMAPSIMKSL MDHTIPEV |
| Tag: | His-tag |
| Predicted MW: | 34.6 kDa |
| Concentration: | lot specific |
| Purity: | >90% |
| Buffer: | Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl Buffer (pH 8.0) containing 100 mM NaCl, 40% Glycerol |
| Preparation: | Liquid purified protein |
| Protein Description: | Recombinant human SDCBP protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography. |
| Storage: | Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| RefSeq: | <u>NP_001007068</u> |
| Locus ID: | 6386 |
| UniProt ID: | <u>O00560</u> , <u>A0A024R7Z5</u> |
| Cytogenetics: | 8q12.1 |
| Synonyms: | MDA-9; MDA9; ST1; SYCL; TACIP18 |



[View online »](#)

Summary:

The protein encoded by this gene was initially identified as a molecule linking syndecan-mediated signaling to the cytoskeleton. The syntenin protein contains tandemly repeated PDZ domains that bind the cytoplasmic, C-terminal domains of a variety of transmembrane proteins. This protein may also affect cytoskeletal-membrane organization, cell adhesion, protein trafficking, and the activation of transcription factors. The protein is primarily localized to membrane-associated adherens junctions and focal adhesions but is also found at the endoplasmic reticulum and nucleus. Alternative splicing results in multiple transcript variants encoding different isoforms. Related pseudogenes have been identified on multiple chromosomes. [provided by RefSeq, Jan 2017]

Protein Families:

Druggable Genome, Transmembrane

Product images: