

Product datasheet for **AR09790PU-L**

PSMB5 (60-263, His-tag) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	PSMB5 (60-263, His-tag) human recombinant protein, 0.25 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	<u>MRGSHHHHHH</u> <u>GMASMTGGQQ</u> <u>MGRDLYDDDD</u> <u>KDRWGSM</u> <u>TTT</u> LAFKFRHGVI VAADSRATAG AYIASQTVKK VIEINPYLLG TMAGGAADCS FWERLLARQC RIYELRNKER ISVAAASKLL ANMVYQYKGM GLSMGMTICG WDKRGPGLYY VDSEGNRISG ATFSVSGSV YAYGVMDRGY SYDLEVEQAY DLARRAIYQA TYRDAYSGBA VNLYHVREDG WIRVSSDNVA DLHEKYSGST P
Tag:	His-tag
Predicted MW:	26.7 kDa
Concentration:	lot specific
Purity:	>90%
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.5) containing 50% glycerol, 1 mM DTT, 0.15M NaCl
Preparation:	Liquid purified protein
Protein Description:	Recombinant human PSMB5 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>NP_001124197</u>
Locus ID:	5693
UniProt ID:	<u>P28074</u> , <u>A0A140VJS7</u>
Cytogenetics:	14q11.2
Synonyms:	LMPX; MB1; X



[View online »](#)

Summary:

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit in the proteasome. This catalytic subunit is not present in the immunoproteasome and is replaced by catalytic subunit 3i (proteasome beta 8 subunit). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2009]

Protein Families:

Protease

Protein Pathways:

Proteasome

Product images: