

# **Product datasheet for AR09793PU-S**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MIP4 / CCL18 (22-89, His-tag) Human Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** MIP4 / CCL18 (22-89, His-tag) human recombinant protein, 10 μg

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** MGSSHHHHHH SSGLVPRGSH MGSHMQVGTN KELCCLVYTS WQIPQKFIVD YSETSPQCPK

or AA Sequence: PGVILLTKRG RQICADPNKK WVQKYISDLK LNA

Tag: His-tag
Predicted MW: 10.4 kDa

Concentration: lot specific
Purity: >95%

**Buffer:** Presentation State: Purified

State: Liquid purified peptide

Buffer System: 10 mM Sodium Citrate pH 3.5, 10% Glycerol

**Endotoxin:** < 1.0 EU per 1 μg of protein (determined by LAL method )

**Preparation:** Liquid purified peptide

**Protein Description:** Recombinant human CCL18, fused to His-tag at N-terminus, was expressed in E.coli and

purified by using conventional chromatography techniques.

**Storage:** Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

RefSeg: NP 002979

 Locus ID:
 6362

 UniProt ID:
 P55774

 Cytogenetics:
 17q12

Synonyms: C-C motif chemokine 18, AMAC1, DCCK1, MIP-4, PARC, Small-inducible cytokine A18, SCYA18





**Summary:** 

This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for naive T cells, CD4+ and CD8+ T cells and nonactivated lymphocytes, but not for monocytes or granulocytes. This chemokine attracts naive T lymphocytes toward dendritic cells and activated macrophages in lymph nodes. It may play a role in both humoral and cell-mediated immunity responses. [provided by RefSeq, Sep 2014]

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

### **Product images:**

