

Product datasheet for AR31199PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD274 / PDL1 (Fc portion) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: CD274 / PDL1 (Fc portion) human recombinant protein, 20 μg

Species: Human
Expression Host: CHO

Expression cDNA Clone <u>Monomer</u>:

or AA Sequence: FTVTVPKDLY VVEYGSNMTI ECKFPVEKQL DLAALIVYWE MEDKNIIQFV HGEEDLKVQH

SSYRQRARLL KDQLSLGNAA LQITDVKLQD AGVYRCMISY GGADYKRITV KVNAPYNKIN

QRILVVDPVT SEHELTCQAE GYPKAEVIWT SSDHQVLSGK TTTTNSKREE KLFNVTSTLR INTTTNEIFY

CTFRRLDPEE NHTAELVIPE LPLAHPPNER GGPKSCDKTH TCPPCPAPEL LGGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV HNAKTKPREE QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKALPAPIEK TISKAKGQPR EPQVYTLPPS RDELTKNQVS LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF FLYSKLTVDK SRWQQGNVFS

CSVMHEALHN HYTQKSLSLS PGK

Predicted MW: 102.6 kDa

Purity: >95% pure by SDS-PAGE and HPLC analyses

Buffer: Presentation State: Purified

State: Lyophilized (sterile filtered) purified protein

Buffer System: 10 mM Sodium Phosphate + 25 mM NaCl, pH 7.6

Bioactivity: Biological: Determined by its ability to induce adhesion in T-cell enriched PBMC cultures. The

ED50 for this effect is 1.2-2.0 μ g/ml.

Endotoxin: $< 0.1 \text{ ng per } \mu\text{g (1EU/}\mu\text{g)}$

Reconstitution Method: Restore in Water to a concentration of 0.1-1.0 mg/ml.

This solution can then be diluted into other aqueous buffers and stored at 2-8°C for 1 week

or -20°C for future use.

Preparation: Lyophilized (sterile filtered) purified protein





CD274 / PDL1 (Fc portion) Human Protein - AR31199PU-S

Protein Description: CHO cell-derived Recombinant Human PD-L1 Fc is a glycosylated, disulfide-linked homodimer

of 906 amino acid residues whose monomer consists of the 220-amino-acid length

extracellular portion of PD-L1 fused to the

231-amino-acid length Fc portion of human IgG1 by two glycines.

The calculated molecular weight of CHO cell-derived Recombinant Human PD-L1 Fc is 102.6

kDa, however, due to glycosylation, it migrates at an apparent molecular weight of approximately 160-170 kDa by SDS-PAGE analysis under non-reducing conditions.

Note: Centrifuge the vial prior to opening!

Storage: Store lyophilized at 2-8°C for 6 months or at -20°C long term.

After reconstitution store the antibody undiluted at 2-8°C for one month

or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001254635

Locus ID: 29126

UniProt ID: Q9NZQ7, Q0GN75

Cytogenetics: 9p24.1

Synonyms: B7-H; B7H1; hPD-L1; PD-L1; PDCD1LG1; PDL1

Summary: This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic

and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq,

Sep 2015]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs)