

Product datasheet for AR31203SU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Helicobacter pylori Protein

Product data:

Product Type: Native Proteins

Description: Helicobacter pylori protein, 0.1 ml

Protein Source: Cell culture

Purity: None, Native Lysate

Buffer: Presentation State: Lysate

State: Liquid Lysate

Buffer System: Phosphate Buffered Saline, pH 7.4

Preservative: 0.09% Sodium Azide

Preparation: Liquid Lysate

Applications: Suitable for use as a positive control in Lateral Flow assays and when spiked into stool

samples.

Protein Description: H. pylori Helicobacter pylori (H. pylori).

Note: Inactivation: Binary Ethylenimine + Formaldehyde Inactivation + Heat treatment at 60°C for

30 minutes. Confirmed by absence of colonies after 72 hours.

Storage: Upon receipt, store (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Synonyms: H. pylori

Summary: The spiral shaped bacterium Helicobacter pylori is strongly associated with inflammation of

the stomach and is also implicated in the development of gastric malignancy. H. pylori is known to cause peptic ulcers and chronic gastritis in human. It is associated with duodenal ulcers and may be involved in development of adenocarcimona and low-grade lymphoma of mucosa associated lymphoid tissue in the stomach. More recently this bacterium has also been implicated with a number of vascular disorders including heart disease. It is not clear how H. pylori is transmitted or why some patients become symptomatic while others do not. The bacteria are most likely spread from person to person through fecal-to-oral or oral-to-oral routes. Possible environmental reservoirs include contaminated water sources.

Protein Families: Suitable for use as a positive control in Lateral Flow assays and when spiked into stool

samples.

