

Product datasheet for AR39087PU-N

UBE2D2 (1-147, His-tag) Human Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Recombinant Proteins |
|--|--|
| Description: | UBE2D2 (1-147, His-tag) human recombinant protein, 0.1 mg |
| Species: | Human |
| Expression Host: | E. coli |
| Expression cDNA Clone or AA Sequence: | <u>MGSSHHHHHH SSGLVPRGSH</u> MALKRIHKEL NDLARDPPAQ CSAGPVGDDM FHWQATIMGP NDSPYQGGVF FLTIHFPTDY PFKPPKVAFT TRIYHPNINS NGSICLDILR SQWSPALTIS KVLLSICSLL CDPNPDDPLV PEIARIYKTD REKYNRIARE WTQKYAM |
| Tag: | His-tag |
| Predicted MW: | 18.9 kDa |
| Concentration: | lot specific |
| Purity: | >95% |
| Buffer: | Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 10% glycerol, 1 mM DTT, 0.1M NaCl |
| Preparation: | Liquid purified protein |
| Protein Description: | Recombinant human UBE2D2 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques. |
| Storage: | Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| RefSeq: | <u>NP 003330</u> |
| Locus ID: | 7322 |
| UniProt ID: | <u>P62837</u> |
| Cytogenetics: | 5q31.2 |
| Synonyms: | E2(17)KB2; PUBC1; UBC4; UBC4/5; UBCH4; UBCH5B |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Serigene UBE2D2 (1-147, His-tag) Human Protein – AR39087PU-N

Summary:Regulated degradation of misfolded, damaged or short-lived proteins in eukaryotes occurs
via the ubiquitin (Ub)-proteasome system (UPS). An integral part of the UPS system is the
ubiquitination of target proteins and covalent linkage of Ub-containing proteins to form
polymeric chains, marking them as targets for 26S proteasome-mediated degradation.
Ubiquitination of proteins is mediated by a cascade of enzymes which includes E1 (ubiquitin
activating), E2 (ubiquitin conjugating), and E3 (ubiquitin ligases) enzymes. This gene encodes a
member of the E2 enzyme family. Substrates of this enzyme include the tumor suppressor
protein p53 and peroxisomal biogenesis factor 5 (PEX5). Alternative splicing results in
multiple transcript variants of this gene. [provided by RefSeq, May 2013]

Protein Pathways: Ubiquitin mediated proteolysis

Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US