

Product datasheet for **AR39131PU-N**

FGF acidic / FGF1 (16-155, His-tag) Mouse Protein

Product data:

Product Type:	Recombinant Proteins
Description:	FGF acidic / FGF1 (16-155, His-tag) mouse protein, 0.1 mg
Species:	Mouse
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MGSSHHHHHH SSGLVPRGSH MFNLPLGNYK KPKLLYCSNG GHFLRILPDG TVDGTRDRSD QHIQLQLSAE SAGEVYIKGT ETGQYLAMDT EGLLYGSQTP NEECLFLERL EENHYNTYTS KKHAENWFV GLKKNNGSCKR GPRTHYGQKA ILFLPLPVSS D
Tag:	His-tag
Concentration:	lot specific
Purity:	>90%
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 1 mM DTT, 30% glycerol, 0.1M NaCl
Endotoxin:	< 1.0 EU per 1 µg of protein (determined by LAL method)
Preparation:	Liquid purified protein
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	NP_034327
Locus ID:	14164
UniProt ID:	P61148 , Q6ZWS1
Cytogenetics:	18 20.74 cM
Synonyms:	Dffrx; Fam; Fgf-1; Fgfa



[View online »](#)

Summary:

Plays an important role in the regulation of cell survival, cell division, angiogenesis, cell differentiation and cell migration. Functions as potent mitogen in vitro. Acts as a ligand for FGFR1 and integrins. Binds to FGFR1 in the presence of heparin leading to FGFR1 dimerization and activation via sequential autophosphorylation on tyrosine residues which act as docking sites for interacting proteins, leading to the activation of several signaling cascades. Binds to integrin ITGAV:ITGB3. Its binding to integrin, subsequent ternary complex formation with integrin and FGFR1, and the recruitment of PTPN11 to the complex are essential for FGF1 signaling. Induces the phosphorylation and activation of FGFR1, FRS2, MAPK3/ERK1, MAPK1/ERK2 and AKT1. Can induce angiogenesis.[UniProtKB/Swiss-Prot Function]

Product images: