

# **Product datasheet for AR50029PU-S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TCEB2 (1-118) (recombinant) Human Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** TCEB2 (1-118) (recombinant) human recombinant protein, 0.1 mg

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** MDVFLMIRRH KTTIFTDAKE SSTVFELKRI VEGILKRPPD EQRLYKDDQL LDDGKTLGEC or AA Sequence: GFTSQTARPQ APATVGLAFR ADDTFEALCI EPFSSPPELP DVMKPQDSGS SANEQAVQ

Predicted MW: 13.1 kDa

Concentration: lot specific

Purity: >95% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 1mMDTT, 0.1 mM PMSF, 20%

glycerol

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant TCEB2 was expressed as insoluble protein aggregate in E.coli and purified by

conventional chromatography, after refolding of the isolated inclusion bodies in a

renaturation buffer.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

RefSeq: NP 009039

**Locus ID:** 6923

 UniProt ID:
 Q15370

 Cytogenetics:
 16p13.3

Synonyms: SIII; TCEB2





**Summary:** 

This gene encodes the protein elongin B, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. Pseudogenes have been identified on chromosomes 11 and 13. [provided by RefSeq, Aug 2008]

**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Pathways in cancer, Renal cell carcinoma, Ubiquitin mediated proteolysis

## **Product images:**

