

Product datasheet for **AR50346PU-N**

CD137 / TNFRSF9 (18-186, His-tag) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	CD137 / TNFRSF9 (18-186, His-tag) human recombinant protein, 0.5 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MGSSHHHHHH SSGLVPRGSH MGSMFERTRS LQDPCSNCPA GTFCDNNRNQ ICSPCPPNSF SSAGGQRTCD ICRQCKGVFR TRKECSSTSN AECDCTPGFH CLGAGCSMCE QDCKQGQELT KKGCKDCCFG TFNDQKRGIC RPWTNCSLDG KSVLVNGTKE RDVVCGPSPA DLSPGASSVT PPAPAREPGH SPQ
Tag:	His-tag
Predicted MW:	20 kDa
Concentration:	lot specific
Purity:	>90% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.1M NaCl, 10% glycerol, 1 mM DTT
Preparation:	Liquid purified protein
Protein Description:	Recombinant human TNFRSF9 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	NP_001552
Locus ID:	3604
UniProt ID:	Q07011
Cytogenetics:	1p36.23
Synonyms:	4-1BB; CD137; CDw137; ILA



[View online »](#)

Summary:

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contributes to the clonal expansion, survival, and development of T cells. It can also induce proliferation in peripheral monocytes, enhance T cell apoptosis induced by TCR/CD3 triggered activation, and regulate CD28 co-stimulation to promote Th1 cell responses. The expression of this receptor is induced by lymphocyte activation. TRAF adaptor proteins have been shown to bind to this receptor and transduce the signals leading to activation of NF-kappaB. [provided by RefSeq, Jul 2008]

Protein Families:

Druggable Genome, Transmembrane

Protein Pathways:

Cytokine-cytokine receptor interaction

Product images: