

Product datasheet for AR50366PU-S

Bcl-2-like 5 (1-152, His-tag) Human Protein

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

Product data:

Product Type: Recombinant Proteins

Description: Bcl-2-like 5 (1-152, His-tag) human recombinant protein, 20 μg

Species: Human
Expression Host: E. coli

Expression cDNA Clone MGSSHHHHHH SSGLVPRGSH MGSHMTDCEF GYIYRLAQDY LQYVLQIPQP GSGPSKTSRV

or AA Sequence: LQKVAFSVQK EVEKNLKSCL DNVNVVSVDT ARTLFNQVME KEFEDDIINW GRIVTIFAFE GILIKKLLRQ

QIAPDVDTYK EISYFVAEFI MNNTGEWIRQ NGGWENGFVK KFEPKS

Tag: His-tag
Predicted MW: 20.2 kDa
Concentration: lot specific

Purity: >95% by SDS - PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 10% glycerol

Preparation: Liquid purified protein

Protein Description: Recombinant human BCL2A1 protein, fused to His-tag at N-terminus, was expressed in E.coli

and purified by using conventional chromatography.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001108207

Locus ID: 597

UniProt ID: Q16548

Cytogenetics: 15q25.1

Synonyms: ACC-1; ACC-2; ACC1; ACC2; BCL2L5; BFL1; GRS; HBPA1





Summary:

This gene encodes a member of the BCL-2 protein family. The proteins of this family form hetero- or homodimers and act as anti- and pro-apoptotic regulators that are involved in a wide variety of cellular activities such as embryonic development, homeostasis and tumorigenesis. The protein encoded by this gene is able to reduce the release of pro-apoptotic cytochrome c from mitochondria and block caspase activation. This gene is a direct transcription target of NF-kappa B in response to inflammatory mediators, and is upregulated by different extracellular signals, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), CD40, phorbol ester and inflammatory cytokine TNF and IL-1, which suggests a cytoprotective function that is essential for lymphocyte activation as well as cell survival. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome
Protein Pathways: Metabolic pathways

Product images:

