

Product datasheet for AR50396PU-S

RGS1 (1-209, His-tag) Human Protein

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Recombinant Proteins
Description:	RGS1 (1-209, His-tag) human recombinant protein, 0.1 mg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MGSSHHHHHH SSGLVPRGSH MGSHMRAAAI STPKLDKMPG MFFSANPKEL KGTTHSLLDD KMQKRRPKTF GMDMKAYLRS MIPHLESGMK SSKSKDVLSA AEVMQWSQSL EKLLANQTGQ NVFGSFLKSE FSEENIEFWL ACEDYKKTES DLLPCKAEEI YKAFVHSDAA KQINIDFRTR ESTAKKIKAP TPTCFDEAQK VIYTLMEKDS YPRFLKSDIY LNLLNDLQAN SLK
Tag:	His-tag
Predicted MW:	26 kDa
Concentration:	lot specific
Purity:	>90% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 10% glycerol, 1 mM DTT
Preparation:	Liquid purified protein
Protein Description:	Recombinant human RGS1 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>NP 002913</u>
Locus ID:	5996
UniProt ID:	<u>Q08116</u>
Cytogenetics:	1q31.2
Synonyms:	1R20; BL34; HEL-S-87; IER1; IR20




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE RGS1 (1-209, His-tag) Human Protein – AR50396PU-S

Summary:

This gene encodes a member of the regulator of G-protein signalling family. This protein is located on the cytosolic side of the plasma membrane and contains a conserved, 120 amino acid motif called the RGS domain. The protein attenuates the signalling activity of G-proteins by binding to activated, GTP-bound G alpha subunits and acting as a GTPase activating protein (GAP), increasing the rate of conversion of the GTP to GDP. This hydrolysis allows the G alpha subunits to bind G beta/gamma subunit heterodimers, forming inactive G-protein heterotrimers, thereby terminating the signal. [provided by RefSeq, Jul 2008]

Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US