

Product datasheet for AR50503PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

KLRC2 / CD159c (94-231, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: KLRC2 / CD159c (94-231, His-tag) human recombinant protein, 0.25 mg

Species: Human
Expression Host: E. coli

Expression cDNA Clone MGSSHHHHHH SSGLVPRGSH MGSMIPFLEQ NNFSPNTRTQ KARHCGHCPE EWITYSNSCY

or AA Sequence: YIGKERRTWE ESLLACTSKN SSLLSIDNEE EMKFLASILP SSWIGVFRNS SHHPWVTING LAFKHKIKDS

DNAELNCAVL QVNRLKSAQC GSSMIYHCKH KL

Tag: His-tag
Predicted MW: 18.4 kDa
Concentration: lot specific

Purity: >85% by SDS - PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.4M urea, 10% glycerol

Preparation: Liquid purified protein

Protein Description: Recombinant human KLRC2 protein, fused to His-tag at N-terminus, was expressed in E.coli.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: <u>NP 002251</u>

 Locus ID:
 3822

 UniProt ID:
 P26717

 Cytogenetics:
 12p13.2

Synonyms: CD159c; NKG2-C; NKG2C





Summary:

Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The group, designated KLRC (NKG2) are expressed primarily in natural killer (NK) cells and encodes a family of transmembrane proteins characterized by a type II membrane orientation (extracellular C terminus) and the presence of a C-type lectin domain. The KLRC (NKG2) gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed on NK cells. KLRC2 alternative splice variants have been described but their full-length nature has not been determined. [provided by RefSeq, Jul 2008]

Protein Families:

Transmembrane

Protein Pathways:

Antigen processing and presentation, Natural killer cell mediated cytotoxicity

Product images:

