OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for AR50571PU-S

MBD3 (1-291, His-tag) Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	MBD3 (1-291, His-tag) human recombinant protein, 50 µg
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	MGSSHHHHHH SSGLVPRGSH MGSMERKRWE CPALPQGWER EEVPRRSGLS AGHRDVFYYS PSGKKFRSKP QLARYLGGSM DLSTFDFRTG KMLMSKMNKS RQRVRYDSSN QVKGKPDLNT ALPVRQTASI FKQPVTKITN HPSNKVKSDP QKAVDQPRQL FWEKKLSGLN AFDIAEELVK TMDLPKGLQG VGPGCTDETL LSAIASALHT STMPITGQLS AAVEKNPGVW LNTTQPLCKA FMVTDEDIRK QEELVQQVRK RLEEALMADM LAHVEELARD GEAPLDKACA EDDDEEDEEE EEEEPDPDPE MEHV
Tag:	His-tag
Predicted MW:	35.2 kDa
Concentration:	lot specific
Purity:	>85% by SDS - PAGE
Buffer:	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.2M NaCl, 40% glycerol, 1 mM DTT, 1 mM EDTA
Preparation:	Liquid purified protein
Protein Description:	Recombinant human MBD3 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Storage:	Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	NP_001268382
Locus ID:	53615
UniProt ID:	O95983
Cytogenetics:	19p13.3

View online »

Summary: DNA methylation is the major modification of eukaryotic genomes and plays an essential role in mammalian development. This gene belongs to a family of nuclear proteins which are characterized by the presence of a methyl-CpG binding domain (MBD). The encoded protein is a subunit of the NuRD, a multisubunit complex containing nucleosome remodeling and histone deacetylase activities. Unlike the other family members, the encoded protein is not capable of binding to methylated DNA. The protein mediates the association of metastasis-associated protein 2 with the core histone deacetylase complex. Alternative splicing results in multiple transcript variants of this gene. [provided by RefSeq, Jul 2013]

Protein Families: Druggable Genome, Transcription Factors

Product images:



15% SDS-PAGE (3ug)