

## Product datasheet for AR50663PU-N

## OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PGPEP1 (1-209, His-tag) Human Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** PGPEP1 (1-209, His-tag) human recombinant protein, 0.5 mg

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

MGSSHHHHHH SSGLVPRGSH MGSMEQPRKA VVVTGFGPFG EHTVNASWIA VQELEKLGLG DSVDLHVYEI PVEYQTVQRL IPALWEKHSP QLVVHVGVSG MATTVTLEKC GHNKGYKGLD NCRFCPGSQC CVEDGPESID SIIDMDAVCK RVTTLGLDVS VTISQDAGRY LCDFTYYTSL

YQSHGRSAFV HVPPLGKPYN ADQLGRALRA IIEEMLDLLE QSEGKINYCH KH

Tag: His-tag
Predicted MW: 25.5 kDa
Concentration: lot specific

Purity: >95% by SDS - PAGE

**Buffer:** Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 20% glycerol, 1 mM

DTT

**Preparation:** Liquid purified protein

**Protein Description:** Recombinant human PGPEP1 protein, fused to His-tag at N-terminus, was expressed in E.coli

and purified by using conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**RefSeq:** <u>NP 001287856</u>

**Locus ID:** 54858

UniProt ID: Q9NX|5, U3KQ24

Cytogenetics: 19p13.11

**Synonyms:** PAP-I; Pcp; PGI; PGP-I; PGPI





**Summary:** 

The gene encodes a cysteine protease and member of the peptidase C15 family of proteins. The encoded protein cleaves amino terminal pyroglutamate residues from protein substrates including thyrotropin-releasing hormone and other neuropeptides. Expression of this gene may be downregulated in colorectal cancer, while activity of the encoded protein may be negatively correlated with cancer progression in colorectal cancer patients. Activity of the encoded protease may also be altered in other disease states including in liver cirrhosis, which is associated with reduced protease activity, and in necrozoospermia, which is associated with elevated protease activity. [provided by RefSeq, Jul 2016]

**Protein Families:** 

Druggable Genome, Protease

## **Product images:**

