

Product datasheet for AR50681PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Bcl-2-like 10 (1-172, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: Bcl-2-like 10 (1-172, His-tag) human recombinant protein, 0.5 mg

Species: Human E. coli **Expression Host:**

Expression cDNA Clone

MGSSHHHHHH SSGLVPRGSH MGSMVDQLRE RTTMADPLRE RTELLLADYL GYCAREPGTP EPAPSTPEAA VLRSAAARLR QIHRSFFSAY LGYPGNRFEL VALMADSVLS DSPGPTWGRV or AA Sequence:

VTLVTFAGTL LERGPLVTAR WKKWGFQPRL KEQEGDVARD CQRLVALLSS RLMGQHRAWL

QAQGGWDGFC HFFRT

Tag: His-tag Predicted MW: 21.8 kDa **Concentration:** lot specific

>90% by SDS - PAGE **Purity:**

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 20% glycerol, 1 mM

DTT

Preparation: Liquid purified protein

Protein Description: Recombinant human BCL2L10 protein, fused to His-tag at N-terminus, was expressed in E.coli

and purified by using conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 001293097

Locus ID: 10017

UniProt ID: Q9HD36, H0YMD5

Cytogenetics: 15q21.2

Synonyms: BCL-B; bcl2-L-10; Boo; Diva





Summary:

The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The protein encoded by this gene contains conserved BH4, BH1 and BH2 domains. This protein can interact with other members of BCL-2 protein family including BCL2, BCL2L1/BCL-X(L), and BAX. Overexpression of this gene has been shown to suppress cell apoptosis possibly through the prevention of cytochrome C release from the mitochondria, and thus activating caspase-3 activation. The mouse counterpart of this protein is found to interact with Apaf1 and forms a protein complex with Caspase 9, which suggests the involvement of this protein in APAF1 and CASPASE 9 related apoptotic pathway. [provided by RefSeq, Jul 2008]

Protein Families:

Druggable Genome, Transmembrane

Product images:

