

Product datasheet for AR50751PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hemoglobin F / HBG2 (1-147, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: Hemoglobin F / HBG2 (1-147, His-tag) human recombinant protein, 0.5 mg

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

MGSSHHHHHH SSGLVPRGSH MGSMGHFTEE DKATITSLWG KVNVEDAGGE TLGRLLVVYP WTQRFFDSFG NLSSASAIMG NPKVKAHGKK VLTSLGDAIK HLDDLKGTFA QLSELHCDKL

VIQKEFDSFG NESSASAINIG NPKVKAHGKK VEISEGDAIK HEDDEKGIFA QESEEHCDI

HVDPENFKLL GNVLVTVLAI HFGKEFTPEV QASWQKMVTG VASALSSRYH

Tag:His-tagPredicted MW:18.5 kDaConcentration:lot specific

Purity: >90% by SDS - PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 0.1M NaCl, 20% glycerol, 1 mM DTT

Preparation: Liquid purified protein

Protein Description: Recombinant human HBG2 protein, fused to His-tag at N-terminus, was expressed in E.coli

and purified by using conventional chromatography techniques.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer. Avoid

repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: NP 000175

Locus ID: 3048

UniProt ID: <u>P69892</u>, <u>D9YZU9</u>

Cytogenetics: 11p15.4

Synonyms: HBG-T1; TNCY





Summary:

The gamma globin genes (HBG1 and HBG2) are normally expressed in the fetal liver, spleen and bone marrow. Two gamma chains together with two alpha chains constitute fetal hemoglobin (HbF) which is normally replaced by adult hemoglobin (HbA) at birth. In some beta-thalassemias and related conditions, gamma chain production continues into adulthood. The two types of gamma chains differ at residue 136 where glycine is found in the G-gamma product (HBG2) and alanine is found in the A-gamma product (HBG1). The former is predominant at birth. The order of the genes in the beta-globin cluster is: 5'- epsilon -- gamma-G -- gamma-A -- delta -- beta--3'. [provided by RefSeq, Jul 2008]

Product images:

